

Correlation functions for 1d interacting fermions with spin-orbit coupling

Aníbal Iucci*

*Instituto de Física La Plata. Departamento de Física, Facultad de Ciencias Exactas,
Universidad Nacional de La Plata. CC 67, 1900 La Plata, Argentina.
Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina.*

We compute correlation functions for one-dimensional electron systems which spin and charge degrees of freedom are coupled through spin-orbit coupling. Charge density waves, spin density waves, singlet- triplet- superconducting fluctuations are studied. We show that the spin-orbit interaction modify the exponents and the phase diagram of the system, changing the dominant fluctuations and making new susceptibilities diverge for low temperature.

PACS numbers: 71.10.Pm

In past two decades there have been an intense effort in studying quasi-one-dimensional electron systems (Q1DES). This interest has its origin in the simplicity of the models which describe them, and at the same time, in the possibility of making contact with experiments. Examples of these Q1DES of recent construction are carbon nanotubes¹, conducting polymers² and semiconductor heterostructures³. From the theoretical point of view the simplest formulation of a Q1DES is given by the Tomonaga-Luttinger^{4,5} model which describes the major qualitative features of the interacting Q1DES such as the spin-charge separation and the non universal exponents in the decay law of correlation functions.

In realistic situations the electrons are moving in electric fields inside the materials: microscopic and macroscopic ones, the latter responsible of confining the electrons to a reduced region of space. As a consequence it appears a magnetic field in the rest frame of the electron which couples with its intrinsic magnetic moment and breaks the spin-rotation SU(2) symmetry. This is known as spin-orbit (SO) interaction (or spin-orbit coupling). Despite its relativistic origin, this interaction has an important effect on existing two-dimensional electron gases (2DEG) such as GaAs/AlGaAs^{6,7} and InGaAs/InAlAs^{8,9} heterostructures. It is responsible for the modification of their band structure by lifting the spin degeneracy and for positive magnetoresistance effects, known as weak antilocalization¹⁰.

In Q1DES there exists an additional potential, responsible of confining the electrons in a narrow channel patterned in 2DEG heterostructures¹¹. Although as far as we know there is no experimental evidence or measures of the strength of the SO coupling resulting from such a potential, theoretical work indicates that it affects quantitatively the splitting energy behavior as a function of the wave vector k , and produces an asymmetric deformation of each spin branch, i.e. the Fermi velocities take different values for different directions of motion¹².

In Ref. 13 the following Hamiltonian was proposed as a model for a Q1DES with SO coupling

$$H = H_0 + H_{\text{int}} \quad (1)$$

where the noninteracting Hamiltonian is

$$H_0 = v_1 \sum_k \left[(k + k_1) c_{kR\uparrow}^\dagger c_{kR\uparrow} - (k - k_1) c_{kL\downarrow}^\dagger c_{kL\downarrow} \right] + v_2 \sum_k \left[(k + k_2) c_{kR\downarrow}^\dagger c_{kR\downarrow} - (k - k_2) c_{kL\uparrow}^\dagger c_{kL\uparrow} \right]. \quad (2)$$

It consists of a modified Tomonaga-Luttinger model that takes into account the asymmetry in the spectrum for each spin branch making $v_1 \neq v_2$ (and $k_1 \neq k_2$). $c_{krs}^\dagger$ creates a right-going ($r=+1$) or left-going ($r=-1$) electron. The interacting Hamiltonian describes forward scattering electron-electron interactions and has a standard form⁵. Umklapp and backscattering terms are irrelevant if we are far from half filling in the former case, and we restrict to repulsive interactions in the latter.

In this article we compute correlation functions for charge density wave (CDW), spin density wave (SDW) $4k_F$ charge density wave ($4k_F$) and singlet- and triplet-superconductivity (SS and TS) operators for the model presented above. The correlation functions for these operators are well-known in the case of zero spin-orbit coupling^{5,14,15}, including logarithmic correction factors^{16,17} and time and temperature dependence¹⁵. We extend these calculations to the case in which SO interaction are present, and study how the exponents of their algebraic decay are modified. We find interesting modifications of the phase diagram of the system when SO interactions are present. For certain regions of the parameter space, SO coupling changes the dominant fluctuations, and makes new susceptibilities diverge for low temperature.

The Hamiltonian (1) can be studied by the use of bosonization technique^{5,18} as in Ref. 13. For convenience we shall define an average velocity $v_0 = (v_1 + v_2)/2$ and the difference $\delta v = v_2 - v_1$, and the same for the Fermi momentum $k_0 = (k_1 + k_2)/2$ and $\delta k = k_2 - k_1$. If we introduce the usual phase fields $\phi_\rho(\phi_\sigma)$ for charge (spin) degrees of freedom and the dual field $\Pi_\rho(\Pi_\sigma)$ the Hamil-

tonian can be represented as

$$H = \frac{v_\rho}{2} \int dx \left[\frac{1}{K_\rho} (\partial_x \phi_\rho)^2 + K_\rho \Pi_\rho^2 \right] + \frac{v_\sigma}{2} \int dx \left[\frac{1}{K_\sigma} (\partial_x \phi_\sigma)^2 + K_\sigma \Pi_\sigma^2 \right] + \delta v \int dx [(\partial_x \phi_\rho) \Pi_\sigma + (\partial_x \phi_\sigma) \Pi_\rho]. \quad (3)$$

$v_{\rho,\sigma}$ are the propagation velocities of the spin and charge collective modes of the decoupled model ($\delta v = 0$), and $K_{\rho,\sigma}$ are the stiffness constants. The spin-orbit interaction appears as an effect that breaks the spin charge separation as reveals the presence of the third term in the last equation. Nevertheless the Hamiltonian (3) can still be diagonalized in terms of two new phase fields which contains a mixture of spin and charge degrees of freedom. The propagation velocities of these collective modes are

$$v_\pm^2 = \frac{v_\sigma^2 + v_\rho^2}{2} + \delta v^2 \pm \sqrt{\left(\frac{v_\rho^2 - v_\sigma^2}{2}\right)^2 + \delta v^2 \left[v_\sigma^2 + v_\rho^2 + v_\rho v_\sigma \left(\frac{K_\rho}{K_\sigma} + \frac{K_\sigma}{K_\rho}\right)\right]}. \quad (4)$$

As $\delta v \rightarrow 0$, $v_+ \rightarrow \max(v_\rho, v_\sigma)$ and $v_- \rightarrow \min(v_\rho, v_\sigma)$. As δv increases, v_- decreases until it vanishes at the points

$$\delta v_\rho^2 = v_\rho v_\sigma \frac{K_\sigma}{K_\rho} \quad (5)$$

$$\delta v_\sigma^2 = v_\rho v_\sigma \frac{K_\rho}{K_\sigma} \quad (6)$$

At these points, the freezing of the lower bosonic mode is accompanied by a divergence in the charge and spin response functions. The static charge compressibility κ diverges at $\delta v = \delta v_\rho$, and at $\delta v = \delta v_\sigma$ occurs a divergence of the static spin susceptibility χ . They behave as

$$\kappa = \kappa_0 \left[1 - \frac{\delta v}{\delta v_\rho} \right]^{-1}, \quad \kappa_0 = \frac{2K_\rho}{\pi v_\rho} \quad (7)$$

$$\chi = \chi_0 \left[1 - \frac{\delta v}{\delta v_\sigma} \right]^{-1}, \quad \chi_0 = \frac{2K_\sigma}{\pi v_\sigma} \quad (8)$$

where κ_0 and χ_0 are the values of κ and χ in absence of SO coupling. Beyond these points the susceptibilities becomes negatives. This behavior of the static response functions together with the vanishing of the collective modes velocity indicates that the system becomes unstable^{5,19} and undergoes a first order phase

transition¹⁷. For $K_\rho > K_\sigma$, δv_ρ turns out to be lower than δv_σ and as δv grows from the zero value the physical divergence takes place in the charge compressibility. This instability is known as phase separation and has been shown to occur in the extended Hubbard model²⁰ and in the t-J model²¹. In the case that $K_\rho < K_\sigma$, the instability takes place in the spin subsystem and is related to the so called metamagnetic transition, observed for instance in the quasi-one-dimensional compound $\text{Ba}_3\text{Cu}_2\text{O}_4\text{Cl}_2$ ²². It also arises in the phase diagram of the XXZ model with next-to-nearest neighbors²³. In presence of a chemical potential (magnetic field), the region where κ (χ) is negative is associated with the coexistence of two phases with different hole concentration (magnetization). The divergence of κ was found in other models with asymmetric dispersion²⁴.

Let us now focus our attention on the correlation functions. Our interest in this work is to obtain their space-time and temperature $T = 1/\beta$ behavior. The operators for CDW, SDW, $4k_F$, SS and TS fluctuations in their bosonized form are

$$\mathcal{O}_{\text{CDW}} = \frac{2}{\pi a} \cos(2k_0 x + \sqrt{2\pi} \phi_\rho) \cos \sqrt{2\pi} \phi_\sigma \quad (9)$$

$$\mathcal{O}_{4k_F} = \frac{1}{(\pi a)^2} \cos(4k_0 x + \sqrt{8\pi} \phi_\rho) \quad (10)$$

$$\mathcal{O}_{\text{SDW},x} = \frac{2}{\pi a} \cos(2k_0 x + \sqrt{2\pi} \phi_\rho) \cos(\delta k x + \sqrt{2\pi} \theta_\sigma) \quad (11)$$

$$\mathcal{O}_{\text{SDW},y} = \frac{2}{\pi a} \cos(2k_0 x + \sqrt{2\pi} \phi_\rho) \sin(\delta k x + \sqrt{2\pi} \theta_\sigma) \quad (12)$$

$$\mathcal{O}_{\text{SDW},z} = \frac{2}{\pi a} \sin(2k_0 x + \sqrt{2\pi} \phi_\rho) \sin \sqrt{2\pi} \phi_\sigma \quad (13)$$

$$\mathcal{O}_{\text{SS}} = \frac{-i}{\sqrt{2\pi} a} e^{-i\sqrt{2\pi} \theta_\rho} \sin \sqrt{2\pi} \phi_\sigma \quad (14)$$

$$\mathcal{O}_{\text{TS},0} = \frac{1}{\sqrt{2\pi} a} e^{-i\sqrt{2\pi} \theta_\rho} \cos \sqrt{2\pi} \phi_\sigma \quad (15)$$

$$\mathcal{O}_{\text{TS},\pm 1} = \frac{1}{2\pi a} e^{\pm i\delta k x} e^{-i\sqrt{2\pi}(\theta_\rho \pm \theta_\sigma)} \quad (16)$$

where a is a short distance cutoff and θ_ν is related to the conjugated field Π_ν by the relation $\Pi_\lambda = \partial_x \theta_\lambda$.

The correlation functions are defined as

$$R_i(x, \tau; \beta) = \langle \mathcal{T}_\tau \mathcal{O}_i(x, \tau) \mathcal{O}_i^\dagger(0, 0) \rangle. \quad (17)$$

where $\mathcal{T}_\tau$ is the (imaginary) time-ordering operator. These objects were calculated in the path integral framework within the Matsubara imaginary time formalism and the results are:

$$R_{\text{CDW}}(x, \tau; \beta) = R_{\text{SDW},z}(x, \tau; \beta) = \frac{\cos 2k_0 x}{2(\pi a)^2} (z_+ \bar{z}_+)^{-(K_\rho \nu_+^\rho + K_\sigma \nu_+^\sigma)/2} (z_- \bar{z}_-)^{-(K_\rho \nu_-^\rho + K_\sigma \nu_-^\sigma)/2} \left[\left(\frac{\bar{z}_+ z_-}{z_+ \bar{z}_-} \right)^{H \text{sign}(x\tau)} + h.c. \right] \quad (18)$$

$$R_{\text{SDW},xy}(x, \tau; \beta) = \frac{\cos 2k_1 x}{2(\pi a)^2} (z_+ \bar{z}_+)^{-(K_\rho \nu_+^\rho + \mu_+^\sigma/K_\sigma)/2 - \theta_+^\sigma} (z_- \bar{z}_-)^{-(K_\rho \nu_-^\rho + \mu_-^\sigma/K_\sigma)/2 - \theta_-^\sigma} + \frac{\cos 2k_2 x}{2(\pi a)^2} (z_+ \bar{z}_+)^{-(K_\rho \nu_+^\rho + \mu_+^\sigma/K_\sigma)/2 + \theta_+^\sigma} (z_- \bar{z}_-)^{-(K_\rho \nu_-^\rho + \mu_-^\sigma/K_\sigma)/2 + \theta_-^\sigma} \quad (19)$$

$$R_{4k_F}(x, \tau; \beta) = \frac{\cos 4k_0 x}{2(\pi a)^4} (z_+ \bar{z}_+)^{-2K_\rho \nu_+^\rho} (z_- \bar{z}_-)^{-2K_\rho \nu_-^\rho} \quad (20)$$

$$R_{\text{SS}}(x, \tau; \beta) = R_{\text{TS},0}(x, \tau; \beta) = \frac{1}{2(2\pi a)^2} (z_+ \bar{z}_+)^{-(\mu_+^\rho/K_\rho + K_\sigma \nu_+^\sigma)/2 + \theta_+^\rho} (z_- \bar{z}_-)^{-(\mu_-^\rho/K_\rho + K_\sigma \nu_-^\sigma)/2 + \theta_-^\rho} + (\theta_\pm^\rho \rightarrow -\theta_\pm^\rho) \quad (21)$$

$$R_{\text{TS},\pm 1}(x, \tau; \beta) = \frac{e^{\pm i\delta k x}}{(2\pi a)^2} (z_+ \bar{z}_+)^{-(\mu_+^\rho/K_\rho + \mu_+^\sigma/K_\sigma)/2} (z_- \bar{z}_-)^{-(\mu_-^\rho/K_\rho + \mu_-^\sigma/K_\sigma)/2} \left(\frac{\bar{z}_+ z_-}{z_+ \bar{z}_-} \right)^{\pm G \text{sign}(x\tau)} \quad (22)$$

where

$$z_\pm = \frac{\sin \frac{\pi}{v_\pm \beta} (v_\pm |\tau| + \epsilon + ix)}{\sin \frac{\pi \epsilon}{v_\pm \beta}} \quad (23)$$

$$\bar{z}_\pm = \frac{\sin \frac{\pi}{v_\pm \beta} (v_\pm |\tau| + \epsilon - ix)}{\sin \frac{\pi \epsilon}{v_\pm \beta}} \quad (24)$$

and the exponents depend on the the stiffness constants multiplied by the factors that include mode velocities dependence. They are given by

$$\nu_\pm^\lambda = \pm \frac{v_\lambda}{v_\pm} \frac{v_\pm^2 - v_{-\lambda}^2 (1 - \delta v^2 / \delta v_{-\lambda}^2)}{v_\pm^2 - v_-^2} \quad (25)$$

$$\mu_\pm^\lambda = \pm \frac{v_\lambda}{v_\pm} \frac{v_\pm^2 - v_{-\lambda}^2 (1 - \delta v^2 / \delta v_\lambda^2)}{v_\pm^2 - v_-^2} \quad (26)$$

$$\theta_\pm^\lambda = \pm \frac{\delta v}{v_\pm} \frac{v_\pm^2 - (\delta v_\lambda^2 - \delta v^2)}{v_\pm^2 - v_-^2} \quad (27)$$

with $\lambda = \rho, \sigma$, and

$$H = \delta v \frac{K_\rho v_\rho + K_\sigma v_\sigma}{v_+^2 - v_-^2} \quad (28)$$

$$G = \delta v \frac{v_\rho/K_\rho + v_\sigma/K_\sigma}{v_+^2 - v_-^2}. \quad (29)$$

$\nu_\pm^\lambda$ and $\mu_\pm^\lambda$ are positive, and $\theta_\pm^\lambda$, G and H has the same sign as δv .

In the model with zero SO coupling the SU(2) symmetry can be restored by imposing the constraint $K_\sigma = 1$, which emerges naturally if the model under study is the continuum limit of a lattice model with only charge density interactions. In this case this is not possible; the SU(2) symmetry keeps broken even for $K_\sigma = 1$ as reveals the differences in the decays between SDW operators correlation functions in the z direction and in the x, y directions. As in the zero SO case, correlation functions for SDW operators in the z direction and CDW operators are equal, and the same happen with TS,0 and SS operators. This degeneracy is broken by logarithmic corrections that arises if irrelevant backscattering or umklapp terms are included¹⁶.

An interesting point to observe is the appearance of two terms in the SDW, xy correlation functions (Eq. 19) where the modulations have different frequencies and decay with different exponents. As $\theta_\pm^\lambda$ has the same sign as δv (see Eq. 27 and the comment below Eq. 29) for $v_2 > v_1$ ($v_2 < v_1$) the dominant term is the one with frequency k_2 (k_1). In other words the biggest frequency dominates. Also $R_{\text{TS},\pm 1}$ becomes oscillating.

Up to here we have obtained very general formulae for space-time and temperature dependent correlation functions for the model under analysis. We can gain physical insight by observing the algebraic decay of the instantaneous correlation functions at zero temperature, and studying how the exponents get modified from the zero




FIG. 1: Behavior of the exponents α_i' s as a function of δv (in units of v_0). For $v_\rho = 1.2v_0$, $v_\sigma = 0.8v_0$, $K_\rho = 0.6$ and $K_\sigma = 0.85$. For $\delta v \gtrsim 0.16$ SDW, xy fluctuations get dominant, and for $\delta v \gtrsim 0.25$ α_{SS} gets positive, and χ_{SS} divergent for $T \rightarrow 0$.

SO case. The functions behave as

$$R_i(x) \sim |x|^{-2+\alpha_i}. \quad (30)$$

The exponents α_i' s determine the divergence of the corresponding Fourier space susceptibility as $T \rightarrow 0$, $\chi_i(T) \sim T^{-\alpha_i'}$. This makes these instabilities of a completely different nature than the ones described in equations (7) and (8). The expressions obtained for the α_i are

$$\alpha_{CDW} = \alpha_{SDW,z} = 2 - K_\rho \nu^\rho - K_\sigma \nu^\sigma \quad (31)$$

$$\alpha_{SDW,x} = \alpha_{SDW,y} = 2(1 + |\theta^\sigma|) - K_\rho \nu^\rho - \mu^\sigma / K_\sigma \quad (32)$$

$$\alpha_{SS} = \alpha_{TS,0} = 2(1 + |\theta^\rho|) - \mu^\rho / K_\rho - K_\sigma \nu^\sigma \quad (33)$$

$$\alpha_{TS,\pm 1} = 2 - \mu^\rho / K_\rho - \mu^\sigma / K_\sigma. \quad (34)$$

These are the new exponents, which retain the same structure as in the zero SO coupling, but modified by the factors

$$\mu^\lambda = \mu_+^\lambda + \mu_-^\lambda \quad (35)$$

$$\nu^\lambda = \nu_+^\lambda + \nu_-^\lambda \quad (36)$$

$$\theta^\lambda = \theta_+^\lambda + \theta_-^\lambda. \quad (37)$$

When $\delta v \rightarrow 0$, $\theta^\lambda \rightarrow 0$ and $\mu^\lambda, \nu^\lambda \rightarrow 1$, so we reproduce the right results for the zero SO case.

For finite SO coupling, δv appears as a parameter which plays a role in determining the slowest decaying correlation function, and which are the divergent susceptibilities. In Fig. 1 we observe as an example, the behavior of the exponents as a function of δv for $v_\rho = 1.2v_0$, $v_\sigma = 0.8v_0$, $K_\rho = 0.6$ and $K_\sigma = 0.85$. For δv small, CDW fluctuations are dominant, but for $\delta v \gtrsim 0.16v_0$ the SDW, xy correlations decay slower. For small δv , CDW and SDW are the only diverging susceptibilities for $T \rightarrow 0$, but for $\delta v \gtrsim 0.25v_0$, α_{SS} becomes positive,




FIG. 2: Phase diagram in $K_\rho - K_\sigma$ space. The phase in brackets is the subdominant one, which becomes dominant for strong enough SO coupling.




FIG. 3: Phase diagram in $K_\rho - \delta v$ space for $v_\rho = 1.2v_0$, $v_\sigma = 0.8v_0$ and different values of K_σ . $\delta v > \delta v_\sigma$ below the dotted line and metamagnetism occurs.

and χ_{SS} divergent for $T \rightarrow 0$. Calculations of the electron band structure modified by SO coupling show that these values of δv should correspond to typical Q1DES¹².

A careful analysis of the exponents allows us to construct a phase diagram in $K_\rho - K_\sigma$ space (Fig. 2). In each region we indicate the dominant fluctuation for small δv , and in brackets, the dominant one for stronger δv . Other subdominant fluctuations are not indicated. Cross sections of the phase diagram are shown in Fig. 3. In this plot the $K_\rho - \delta v$ space can be observed for $K_\rho < 1$ and different values of K_σ . For small δv CDW fluctuations dominate and for stronger δv the system can be either

in the SDW or in the SS phase depending on the values of K_ρ and K_σ . In the region below the dotted line, $\delta v > \delta v_\sigma$, the static spin susceptibility becomes negative and metamagnetism takes place

In conclusion, we have computed correlation functions for a model of one-dimensional correlated electrons with SO coupling. This coupling destroys the spin charge separation as was shown in Ref. 13, and modifies the exponents of correlations decay. As a consequence the phase diagram gets modified. For strong enough SO coupling, it changes the dominant fluctuation, and makes new susceptibilities diverge for $T \rightarrow 0$. How logarithmic corrections originated in irrelevant backscattering and/or umk-

lapp terms modify these results is an interesting problem, subject of future work.

Acknowledgments

This work was partially supported by the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET) and Universidad Nacional de La Plata (UNLP), Argentina. I am grateful to Carlos Naón for useful comments and discussions, and for encouraging me to write this article.

-
- * Electronic address: iucci@fisica.unlp.edu.ar
- ¹ M. Bockrath, D. Cobden, and J. Lu, A. Rinzler, R. Smalley, L. Balents, and P. Mc-Euen, *Nature* **397**, 598 (1999).
 - ² A. J. Heeger, S. A. Kivelson, J. R. Schrieffer, and W. P. Su, *Rev. Mod. Phys.* **60**, 781 (1988).
 - ³ S. Tarucha, T. Honda, and T. Saku, *Solid State Comm.* **94**, 413 (1995).
 - ⁴ S. Tomonaga, *Prog. Theor. Phys.* **5**, 544 (1950); J. Luttinger, *J. Math. Phys.* **4**, 1154 (1963); D. Mattis and E. Lieb, *J. Math. Phys.* **6**, 304 (1965).
 - ⁵ J. Voit, *Rep. Prog. Phys.* **58**, 977 (1995).
 - ⁶ J. B. Miller, D. M. Zumbuhl, C. M. Marcus, Y. B. Lyanda-Geller, D. Goldhaber-Gordon, K. Campman, A. C. Gosard, *Phys. Rev. Lett.* **90**, 076807 (2003).
 - ⁷ T. Hassenkam, S. Pedersen, K. Baklanov, A. Kristensen, C. B. Sorensen, P. E. Lindelof, F. G. Pikus, and G. E. Pikus, *Phys. Rev. B* **55**, 9298 (1997).
 - ⁸ J. Luo, H. Munekata, F. F. Fang, and P. J. Stiles, *Phys. Rev. B* **41**, 7685 (1990).
 - ⁹ J. Nitta, T. Akazaki, and H. Takayanagi, T. Enoki, *Phys. Rev. Lett.* **78**, 1335 (1997).
 - ¹⁰ G. Bergmann, *Phys. Rep.* **107**, 1 (1984).
 - ¹¹ U. Meirav, M. A. Kastner, M. Heiblum, and S. J. Wind, *Phys. Rev. B* **40**, 5871 (1989).
 - ¹² A. V. Moroz and C. H. W. Barnes, *Phys. Rev. B* **60**, 14272 (1999); *Phys. Rev. B* **61**, R2464 (2000).
 - ¹³ A. V. Moroz, K. V. Samokhin and C. H. W. Barnes, *Phys. Rev. Lett.* **84**, 4164 (2000); *Phys. Rev. B* **62**, 16900 (2000).
 - ¹⁴ J. Solyom, *Adv. Phys.* **28**, 209 (1979).
 - ¹⁵ V. J. Emery, in *Highly Conducting One-Dimensional Solids*, edited by J. T. Devreese *et al.* (Plenum, New York, 1979), p. 327.
 - ¹⁶ T. Giamarchi and H. J. Schulz, *Phys. Rev. B* **39**, 4620 (1989); I. Affleck, D. Gepner, T. Ziman, and H. Schulz, *J. Phys. A* **22**, 511 (1989).
 - ¹⁷ J. Voit, *Phys. Rev. B* **45**, 4027 (1992).
 - ¹⁸ F. D. M. Haldane, *J. Phys. C* **14**, 2585 (1981); J. von Delft and H. Schoeller, *Ann. Phys. (Leipzig)* **7**, 225 (1998).
 - ¹⁹ J. Drut and D. Cabra, *J. Phys.: Condens. Matter* **15**, 1445 (2003).
 - ²⁰ K. Penc and F. Mila, *Phys. Rev. B* **49**, 9670 (1994).
 - ²¹ M. Ogata, M. U. Luchini, S. Sorella, and F. F. Assaad, *Phys. Rev. Lett.* **66**, 2388 (1991).
 - ²² D. Eckert, K. Ruck, M. Wolf, G. Krabbes, and K.-H. Müller, *J. Appl. Phys.* **83**, 7240 (1998).
 - ²³ C. Gerhardt, K.-H. Mütter, and H. Kröger, *Phys. Rev. B* **57**, 11504 (1998).
 - ²⁴ V. I. Fernández, A. Iucci, and C. M. Naón, *Eur. Phys. J. B* **30**, 53 (2002).